



Comprehensive quality assessment of GOME- and IASI-type multi-mission tropospheric ozone data records

Arno Keppens, Jean-Christopher Lambert, D. Hubert, J. Granville, T. Verhoelst, S. Compernelle, J. van Peet, R. van Der A, G. Miles, R. Siddans, et al.

► To cite this version:

Arno Keppens, Jean-Christopher Lambert, D. Hubert, J. Granville, T. Verhoelst, et al.. Comprehensive quality assessment of GOME- and IASI-type multi-mission tropospheric ozone data records. Quadrennial Ozone Symposium 2016, Sep 2016, Edimbourg, United Kingdom. , pp.QOS2016-269-1. insu-01369154

HAL Id: insu-01369154

<https://hal-insu.archives-ouvertes.fr/insu-01369154>

Submitted on 20 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comprehensive quality assessment of GOME- and IASI-type multi-mission tropospheric ozone data records

A. Keppens*, J.-C. Lambert, D. Hubert, J. Granville, T. Verhoelst, S. Compennolle

Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

C. Clerbaux, P.-F. Coheur, G. Miles, R. Siddans, J. van Peet, R. van der A, A. Delcloo, S. Godin-Beekmann, R. Kivi, R. Stübi, C. Zehner**

INTRODUCTION


- Tropospheric ozone plays a key role in air quality and has a significant impact on the radiation budget of the Earth, both directly and indirectly.
- Tropospheric ozone observations are provided by two series of European nadir-viewing ozone profilers:
 - UV-visible spectrometers of the GOME type, launched regularly since 1995: GOME, SCIAMACHY, OMI, GOME-2 on MetOp-A/B/C, upcoming Sentinel-4/5/5p missions
 - Thermal infrared sounders of the IASI type, launched regularly since 2006: IASI on MetOp platforms and IASI-NG on MetOp-SG
- Several tropospheric ozone data products have been improved and harmonised in the context of the ESA's Climate Change Initiative (CCI) on ozone.
- To verify their fitness-for-purpose, we have applied to the Ozone CCI datasets a QA/validation expert system developed over years in the context of ESA's Multi-TASTE and CCI projects, EUMETSAT's O3M-SAF, and the European Commission's GEOmon and QA4ECV.

DATA SELECTION / CO-LOCATION / HARMONISATION

- GOME and GOME-2 nadir UV retrievals by RAL:
 - RAL v2.14
 - 20 fixed retrieval levels (VMR & ND) → 19 layers (PC)
- IASI TIR retrievals by ULB/LATMOS:
 - FORLI v20140922
 - 41 fixed retrieval layers (PC)

Satellite instrument	L2 ozone profile processing entity	Time period
		96 97 98 99 00 01 02 03 04 05 06 07 08 09 10 11 12 13 14 15
GOME	RAL	delivered
SCIAMACHY	RAL	
GOME2A	RAL	expected
GOME2B	RAL	
IASI	ULB/LATMOS	


- 100 km and 3 h co-locations with ozonesonde data at 149 stations from WMO GAW contributing networks like NDACC and SHADOZ



- Mass-conservation regridding before optional AK-based vertical smoothing.
- Tropocolumn from direct summation over partial column profiles
- Two 'tropopause' products are considered:
 - Lapse-rate tropopause (LRT WMO def.) based on ozonesonde T profile
 - Fixed-level cut-off at 6 km (adopted within CCI)

INFORMATION CONTENT STUDY


- From fractional averaging kernel matrices!
- Number of independent layers (DFS) shows meridian and temporal dependence.
- Variations of vertical sensitivity and other information content measures are directly related to slant column density (SCD), affected by solar zenith angle (SZA), latitude...
- Impact of instrument degradation on information content appears clearly.
- Poor sensitivity below tropopause (TP) increases tropocolumn uncertainty.



GROUND-BASED NETWORK VALIDATION

Half-year running median relative difference (thick lines) and 68 % IP spread (thin lines) for LRT-integrated tropospheric ozone column comparison with sonde network data:

- BIAS: GOME(-2A) bias of 10-25 % with stronger outliers, smaller (5-10 % negative) IASI bias due to higher sensitivity below TP
- SMOOTH: Impact of vertical smoothing up to same 25 % order as GOME(-2A) bias
- DRIFT: Pos. for GOME-2A, neg. for GOME, strongest drift for GOME at low latitudes (from 3-year median as green stairs for vertically smoothed data)
- TP-DEF: 6 km fixed cut-off bias (not shown) typically slightly larger, with amplified seasonal bias variation for all instruments




* Contact: arno.keppens@aeronomie.be or see www.esa-ozone-cci.org



aeronomie.be



EUMETSAT



ACKNOWLEDGMENTS

Funding: ESA
Ozone_cci and ProdEx
A3C.

Data: NDACC and GAW
Pls and staff at
stations. Satellite
science and processing
teams.

** Affiliations
Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands
Rutherford Appleton Laboratory (RAL), Chilton, Didcot, UK
Université Libre de Bruxelles (ULB), Brussels, Belgium
LATMOS/RSI, UPMC, UVSQ, CNRS, Paris, France
Royal Meteorological Institute of Belgium (RMI), Brussels, Belgium
Finnish Meteorological Institute (FMI-AK), Sodankylä, Finland
Federal Office of Meteorology and Climatology, Payerne, Switzerland
European Space Agency (ESA/ESRIN), Frascati, Italy

